



University of Insubria

DIPARTIMENTO DI SCIENZA E ALTA TECNOLOGIA  
PhD in Computer Science and Mathematics of Calculus

REPORT OF AN INTRODUCTION TO REGULARITY STRUCTURES

**A Regularity Structure for Rough Volatility**

Teacher: *F. Caravenna*    Student: *M. Tarsia\**    Tutor: *D. Cassani*    Advisor: *E. Mastrogiacomo*

Academic Year 2019–2020

23/02/2020

**Hölder distributions**

Consider  $d \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}$  and  $\mathbb{R}^d = \mathbb{R}_x^d$ ,  $x = (x_1, \dots, x_d)$ , equipped by the Euclidean topology and the Borel  $\sigma$ -algebra  $\mathcal{B}(\mathbb{R}^d)$  with the Lebesgue measure on it. We denote by  $\mathcal{K}_d$  the subclass of  $\mathcal{B}(\mathbb{R}^d)$  consisting of the compact subsets  $K$  of  $\mathbb{R}^d$  and, for  $\rho \in \mathbb{R}_+^* = ]0, \infty[$  and  $x \in \mathbb{R}^d$ , by  $B_\rho(x)$  the open ball in  $\mathbb{R}^d$  of radius  $\rho$  and center  $x$ . Whenever  $x = 0$ , we write  $B_\rho := B_\rho(0)$ .

A *smooth test* or *bump function* on  $\mathbb{R}^d$  is a function  $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$  which is infinitely differentiable and compactly supported. The space of the test functions on  $\mathbb{R}^d$  is denoted by  $\mathcal{D}_d := \mathcal{D}(\mathbb{R}^d) \doteq C_c^\infty(\mathbb{R}^d; \mathbb{R})$  and it's associated to the topology induced by the following uniform convergence notion: given  $\varphi, (\varphi_n)_n$  in  $\mathcal{D}_d$ ,  $\varphi_n \rightarrow \varphi$  in  $\mathcal{D}_d$  as  $n \rightarrow \infty$  if there exists  $K \in \mathcal{K}_d$  with  $\text{supp } \varphi \cup \text{supp } \varphi_n \subseteq K$  for  $n$  large enough and, for any multi-index  $k = (k_1, \dots, k_d) \in \mathbb{N}^d$ ,  $\|D^k \varphi - D^k \varphi_n\|_\infty \rightarrow 0$  as  $n \rightarrow \infty$ , where  $D^k = \partial_{x_1}^{k_1} \dots \partial_{x_d}^{k_d}$  and  $\|\cdot\|_\infty$  is the usual uniform norm. Thereby,  $\mathcal{D}_d$  is a complete and locally convex topological not metrizable (real) vector space satisfying the so-called Heine-Borel property. For  $r \in \mathbb{N}^*$ , we denote

$$\mathcal{D}_{d,r} := \{ \varphi \in \mathcal{D}_d \mid \text{supp } \varphi \subset B_r \text{ and } \|\varphi\|_{C^r} \leq 1 \}$$

where  $\|\cdot\|_{C^r} = \max_{k \in \mathbb{N}^d, |k| \leq r} \|D^k \cdot\|_\infty$  and  $|k| = k_1 + \dots + k_d$ .

A *Schwartz distribution* or *generalized function* on  $\mathbb{R}^d$  is a function  $\eta: \mathcal{D}_d \rightarrow \mathbb{R}$  which is linear and (sequentially) continuous w.r.t. the above topology on  $\mathcal{D}_d$ . The space of the distributions on  $\mathbb{R}^d$  is therefore the dual space of  $\mathcal{D}_d$ , in symbols  $\mathcal{D}'_d := (\mathcal{D}_d)'$ , and as such it's paired with the weak-star topology induced by this pointwise convergence notion: given  $\eta, (\eta_n)_n$  in  $\mathcal{D}'_d$ ,  $\eta_n \rightarrow \eta$  in  $\mathcal{D}'_d$  as  $n \rightarrow \infty$  if, for any  $\varphi \in \mathcal{D}_d$ ,  $\eta_n(\varphi) \rightarrow \eta(\varphi)$  (in  $\mathbb{R}$ ) as  $n \rightarrow \infty$ . Hence,  $\mathcal{D}'_d$  is a locally convex topological not metrizable vector space. For  $\eta \in \mathcal{D}'_d$  and  $\varphi \in \mathcal{D}_d$ , we write  $\langle \eta; \varphi \rangle := \eta(\varphi)$ .

Every function  $f \in L^1_{\text{loc}}(\mathbb{R}^d; \mathbb{R})$  is identified with the distribution (on  $\mathbb{R}^d$ ) defined by  $\varphi \mapsto \int_{\mathbb{R}^d} f \varphi dx$ ,  $\varphi \in \mathcal{D}_d$ , and w.r.t. that standpoint  $\mathcal{D}_d$  results a weak-star dense subset of  $\mathcal{D}'_d$ . Even every Radon (tight) measure  $\mu$  on  $\mathcal{B}(\mathbb{R}^d)$  is viewed as a distribution, that given by  $\varphi \mapsto \int_{\mathbb{R}^d} \varphi \mu(dx)$ ,  $\varphi \in \mathcal{D}_d$ .

For  $\eta \in \mathcal{D}'_d$  and  $k \in \mathbb{N}^d$ , the *(distributional) derivative*  $D^k \eta \in \mathcal{D}'_d$  of order  $k$  of  $\eta$  takes values

$$\langle D^k \eta; \varphi \rangle \doteq (-1)^{|k|} \langle \eta; D^k \varphi \rangle, \quad \varphi \in \mathcal{D}_d.$$

For  $\varphi \in \mathcal{D}_d$ ,  $\lambda \in ]0, 1]$  and  $x \in \mathbb{R}^d$ , the *scaled function*  $\varphi_x^\lambda \in \mathcal{D}_d$  is constructed from  $\varphi$  reducing it by factor  $\lambda$  and centering it at point  $x$  without changing its integral on  $\mathbb{R}^d$ :

$$\varphi_x^\lambda(\cdot) \doteq \lambda^{-d} \varphi(\lambda^{-1}(\cdot - x)).$$

Whenever  $x = 0$ , we write  $\varphi^\lambda := \varphi_0^\lambda$ . Note that, if  $\varphi \in \mathcal{D}_{d,r}$  for  $r \in \mathbb{N}^*$ , then  $\text{supp } \varphi_x^\lambda \subset B_{\lambda r}(x)$  and  $\|\varphi_x^\lambda\|_{C^r} \leq \lambda^{-r}$ . Moreover, for any  $k \in \mathbb{N}^d$ ,  $D^k \varphi_x^\lambda = \lambda^{-|k|} (D^k \varphi)_x^\lambda$ .

\*E-mail: [mtarsia1@uninsubria.it](mailto:mtarsia1@uninsubria.it). Web page: <https://www.uninsubria.it/hpp/marco.tarsia>.

For  $\alpha \in \mathbb{R}_+ = [0, \infty[$ , if  $\alpha \notin \mathbb{N}$  then, leaning on the concept of (locally) Hölder continuous function, the space of  $\alpha$ -Hölder functions on  $\mathbb{R}^d$  is defined as

$$\mathcal{C}_d^\alpha := \mathcal{C}^\alpha(\mathbb{R}^d; \mathbb{R}) \doteq \left\{ f \in C^{[\alpha]}(\mathbb{R}^d; \mathbb{R}) \mid \forall k \in \mathbb{N}^d \text{ with } |k| = [\alpha], D^k f \in \mathcal{C}_{\text{loc}}^{\{\alpha\}}(\mathbb{R}^d; \mathbb{R}) \right\}$$

whereas, for  $\alpha \in \mathbb{R}_- = ]-\infty, 0]$ , if  $\alpha \notin \mathbb{Z}$  then the space of  $\alpha$ -Hölder distributions on  $\mathbb{R}^d$  is defined as

$$\mathcal{C}_d^\alpha := \left\{ \eta \in \mathcal{D}'_d \mid \forall K \in \mathcal{K}_d, \exists C_K \in \mathbb{R}_+^* : \forall \varphi \in \mathcal{D}_{d, [-\alpha]}, \lambda \in ]0, 1] \text{ and } x \in K, |\langle \eta; \varphi_x^\lambda \rangle| \leq C_K \lambda^\alpha \right\}$$

and finally, for  $r \in \mathbb{N}^*$ , the space of Hölder distributions of finite order  $r$  on  $\mathbb{R}^d$  is defined as

$$\mathcal{C}_d^{-r} := \left\{ \eta \in \mathcal{D}'_d \mid r \text{ is the min of } s \in \mathbb{N}^* \text{ s.t., } \forall K \in \mathcal{K}_d, \exists C_K \in \mathbb{R}_+^* : \forall \varphi \in \mathcal{D}_d, |\langle \eta; \varphi \rangle| \leq C_K \|\varphi\|_{C^s} \right\}.$$

Every distribution in  $\mathcal{C}_d^{-r}$  is canonically definable on  $C_c^r(\mathbb{R}^d; \mathbb{R})$  (through a continuity standard argument). Observe that, for any  $\alpha, \beta \in \mathbb{R} \setminus \mathbb{N}$ , if  $\alpha < \beta$  then  $\mathcal{C}_d^\beta \subset \mathcal{C}_d^\alpha$ . Moreover, for any  $k \in \mathbb{N}^d$ ,  $D^k(\mathcal{C}_d^\alpha) \subset \mathcal{C}_d^{\alpha-|k|}$ .

Remember the main issues about the operation of product between distributions, summarized by the Schwartz impossibility theorem, and yet the importance of Hölder distributions regarding that.

As far as  $d = 1$ , we omit the subscript  $d$  from each of the previously defined spaces.

### Regular wavelets

For  $n \in \mathbb{N}$ , we denote by  $\mathbb{R}^d[x; n]$  the class of the real polynomials on  $\mathbb{R}^d = \mathbb{R}_x^d$  of degree  $n$ .

**Theorem** (Daubechies, '88). *Given  $r \in \mathbb{N}^*$ , there exists  $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$  with the following properties.*

1. *The function  $\varphi$  is of class  $C_b^r$  and has compact support.*
2. *For every  $P \in \mathbb{R}^d[x; r]$ , there exists  $\hat{P} \in \mathbb{R}^d[x; r]$  such for which  $P(\cdot) = \sum_{h \in \mathbb{Z}^d} \hat{P}(h) \varphi(\cdot - h)$ .*
3. *For any  $h \in \mathbb{Z}^d$ ,  $\int_{\mathbb{R}^d} \varphi(x) \varphi(x - h) dx = \delta_{h,0}$ .*
4. *There exists a sequence  $(a_h)_{h \in \mathbb{Z}^d}$  in  $\mathbb{R}$  such that, for any  $x \in \mathbb{R}^d$ ,  $2^{-d/2} \varphi(x/2) = \sum_{h \in \mathbb{Z}^d} a_h \varphi(x - h)$ .*

The existence of such a function  $\varphi$  is equivalent to the existence of a wavelet basis of  $L_d^2 = L^2(\mathbb{R}^d; \mathbb{R})$  consisting of  $\|\cdot\|_{L_d^2}$ -orthonormal  $C_b^r$  functions with compact support (proceeding according to the wavelet standard analysis of Meyer, '92). Indeed, given  $r \in \mathbb{N}^*$  and taken  $\varphi$  as in the previous theorem, for  $n \in \mathbb{N}$  we set  $\Lambda^n := 2^{-n} \mathbb{Z}^d$  and, for any  $\psi \in C_c^r(\mathbb{R}^d; \mathbb{R})$  and  $y \in \Lambda^n$ ,  $\psi_y^n(\cdot) := 2^{nd/2} \psi(2^n(\cdot - y))$ . Then there exists  $\Phi \subset C_c^r(\mathbb{R}^d; \mathbb{R})$  with  $|\Phi| = \#\Phi < \infty$  which is orthogonal to  $\mathbb{R}^d[x; r]$  and such that  $\{\varphi_y^0\}_{y \in \mathbb{Z}^d} \cup \{\hat{\varphi}_y^n\}_{\hat{\varphi} \in \Phi, n \in \mathbb{N}, y \in \Lambda^n}$  constitutes an orthonormal basis of  $L_d^2$  (intimately related to  $\mathcal{C}_d^{-r}$ ...).

### Regularity structures

A *regularity structure* is a triple  $\mathcal{T} = (A, T, G)$  made by the following three elements.

- An *index set*  $A$ : a subset of  $\mathbb{R}$  with  $0 \in A$  which is bounded from below and locally finite.
- A *model space*  $T$ : a graded vector space indexed over  $A$  of finite-dimensional vector spaces  $T_\alpha$ ,  $\alpha \in A$ , each of which admits basis of *symbols*  $\{\tau_{\alpha,i}\}_{i \in I_\alpha}$ ,  $|I_\alpha| < \infty$ , i.e.  $T = \bigoplus_{\alpha \in A} T_\alpha = \bigoplus_{\alpha \in A} \langle \tau_{\alpha,i} \mid i \in I_\alpha \rangle$ , where  $T_0 = \langle \mathbf{1} \rangle \cong \mathbb{R}$ . Elements  $\tau_\alpha \in T_\alpha$ ,  $\alpha \in A$ , are said to have *homogeneity* or *degree*  $|\tau_\alpha|$  equal to  $\alpha$ . Given  $\tau \in T$  and  $\alpha \in A$ , we write  $\|\tau\|_\alpha$  for a chosen norm  $\|\cdot\|$  of its component in  $T_\alpha$ .
- A *structure group*  $G$ : a  $\circ$ -group of linear operators  $\Gamma$  acting on  $T$  with  $\Gamma \mathbf{1} = \mathbf{1}$  which satisfy a nilpotency property in the meaning that, for any  $\alpha \in A$  and  $\tau_\alpha \in T_\alpha$ ,

$$\Gamma \tau_\alpha - \tau_\alpha \in T_{<\alpha} \doteq \bigoplus_{\alpha' < \alpha} T_{\alpha'}.$$

A model for  $\mathcal{T}$  on  $\mathbb{R}^d$  is then a pair  $M = (\Pi, \Gamma)$  composed of the following two families.

- A map  $\Gamma = (\Gamma_{x,y})_{x,y \in \mathbb{R}^d}: \mathbb{R}^d \times \mathbb{R}^d \rightarrow G$  such that, for any  $x, y, z \in \mathbb{R}^d$ ,  $\Gamma_{x,y} \circ \Gamma_{y,z} = \Gamma_{x,z}$ .
- A map  $\Pi = (\Pi_x)_{x \in \mathbb{R}^d}: \mathbb{R}^d \rightarrow \mathcal{L}(T; \mathcal{D}'_d)$  such that, for any  $x, y \in \mathbb{R}^d$ ,  $\Pi_x \circ \Gamma_{x,y} = \Pi_y$  and furthermore, letting  $r$  be the smallest integer  $> |\min A|$  then, for any  $\gamma \in \mathbb{R}_+^*$  and  $K \in \mathcal{K}_d$ , there exists  $C = C_{\gamma,K} \in \mathbb{R}_+^*$  such that, by varying  $\alpha \in A$  with  $\alpha \leq \gamma$ ,  $\tau_\alpha \in T_\alpha$ ,  $\varphi \in \mathcal{D}_{d,r}$ ,  $\lambda \in ]0, 1]$ ,  $x, y \in K$  and  $\alpha' < \alpha$ ,

$$|(\Pi_x \tau_\alpha)(\varphi_x^\lambda)| \leq C \lambda^\alpha \|\tau_\alpha\| \quad \text{and} \quad \|\Gamma_{x,y} \tau_\alpha\|_{\alpha'} \leq C |x - y|^{\alpha - \alpha'} \|\tau_\alpha\|.$$

We say that  $\Pi_x$ ,  $x \in \mathbb{R}^d$ , realizes an element of  $T$  as a distribution on  $\mathbb{R}^d$ .

### Skorohod integral

Let  $(\Omega, \mathcal{F}, \mathbf{P})$  be a complete probability space such that there exists a 1D standard Brownian motion  $W = (W(t))_{t \geq 0}$  on it, provided with the filtration  $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$  generated by  $W$ , and fix also  $T \in \mathbb{R}_+^*$ .

For any  $n \in \mathbb{N}^*$ , the  $n$ -fold iterated or multiple Itô integral (w.r.t.  $W$ ) of a symmetric function  $f$  in  $L_{n,T}^2 := L^2([0, T]^n; \mathbb{R})$  is the random variable in  $L^2(\mathbf{P}) := L^2((\Omega, \mathcal{F}, \mathbf{P}); \mathbb{R})$  computed as

$$I_n(f) \doteq \int_{[0, T]^n} f(t_1, \dots, t_n) dW(t_1) \cdots dW(t_n).$$

We remark that  $I_n(f)$  is  $\mathcal{F}_T$ -measurable with  $\mathbf{E}[I_n(f)] = 0$  and  $\mathbf{E}[I_n(f)^2] = n! \|f\|_{L_{n,T}^2}^2$ .

**Theorem** (Wiener-Itô chaos expansion). *Let  $X \in L^2(\mathbf{P})$  be  $\mathcal{F}_T$ -measurable. Then there exists an (essentially) unique sequence  $(f_n)_{n \in \mathbb{N}^*}$  of symmetric functions  $f_n$  in  $L_{n,T}^2$  such that, as limit in  $L^2(\mathbf{P})$ ,*

$$X = \mathbf{E}[X] + \sum_{n=1}^{\infty} I_n(f_n).$$

Furthermore,  $\mathbf{E}[X^2] = \mathbf{E}[X]^2 + \sum_{n=1}^{\infty} n! \|f_n\|_{L_{n,T}^2}^2$ .

Now let  $u = (u(t))_{t=0}^T$  be a 1D (stochastic) process such that, for any  $t \in [0, T]$ ,  $u(t) \in L^2(\mathbf{P})$  is  $\mathcal{F}_T$ -measurable, and therefore let  $(f_n(\cdot, t))_{n \in \mathbb{N}^*}$  be the sequence of symmetric functions  $f_n(\cdot, t)$  in  $L_{n,T}^2$  which determine the Wiener-Itô chaos expansion of  $u(t)$ . Consider, for any  $n \in \mathbb{N}^*$ , the symmetrization  $\tilde{f}_n \in L_{n+1,T}^2$  of  $f_n(\cdot, t)$  as a  $L_{n+1,T}^2$ -function defined on  $[0, T]^n \times [0, T]_t$ : for  $t_1, \dots, t_n, t \in [0, T]$ ,

$$\tilde{f}_n(t_1, \dots, t_n, t) = \frac{1}{n+1} \left\{ f_n(t_1, \dots, t_n, t) + \sum_{j=1}^n f_n(t_1, \dots, t_{j-1}, t, t_{j+1}, \dots, t_n, t_j) \right\}.$$

Let's assume also that  $\mathbf{E}[\int_0^T u^2(t) dt] < \infty$ . Then  $u$  is Skorohod integrable (w.r.t.  $W$ ) if the series

$$\int_0^T u(t) \delta W(t) := \delta(u) \doteq \sum_{n=0}^{\infty} I_{n+1}(\tilde{f}_n)$$

converges in  $L^2(\mathbf{P})$  and, in such a case,  $u \in \text{Dom } \delta$  and  $\delta(u)$  is the Skorohod integral of  $u$  (w.r.t.  $W$ ).

Seeing  $\delta$  as an operator from  $\text{Dom } \delta$  into  $L^2(\mathbf{P})$ , called the divergence operator,  $\delta$  results an unbounded and closed linear operator with, for  $u \in \text{Dom } \delta$ ,  $\mathbf{E}[\delta(u)] = 0$  and  $\mathbf{E}[\delta(u)^2] = \sum_{n=0}^{\infty} (n+1)! \|\tilde{f}_n\|_{L_{n+1,T}^2}^2$ .

**Theorem.** *Let  $u = (u(t))_{t=0}^T$  be a 1D process which is  $(\mathcal{F}_t)_{t=0}^T$ -adapted with  $\mathbf{E}[\int_0^T u^2(t) dt] < \infty$ . Then  $u$  is Skorohod integrable and its Skorohod integral coincides with its Itô integral (w.r.t.  $W$ ):*

$$\delta(u) = \int_0^T u(t) dW(t).$$

Keep in mind that in general, for arbitrary  $u \in \text{Dom } \delta$  and  $\zeta: \Omega \rightarrow \mathbb{R}$  random variable which is  $\mathcal{F}_T$ -measurable and such that  $\zeta u := (\zeta u(t))_{t=0}^T \in \text{Dom } \delta$ , then  $\delta(\zeta u) \neq \zeta \delta(u)$ .

### Wick product

A *rapidly decreasing (smooth) function* on  $\mathbb{R}^d$  is a function  $\varphi \in C^\infty(\mathbb{R}^d; \mathbb{R})$  such that, for any  $n \in \mathbb{N}$  and  $k \in \mathbb{N}^d$ ,  $\|\varphi\|_{n,k} \doteq \sup_{x \in \mathbb{R}^d} |x|^n |D^k \varphi(x)| < \infty$ . The space of the rapidly decreasing functions on  $\mathbb{R}^d$  is the *Schwartz space* (w.r.t.  $d$ ), is denoted by  $\mathcal{S}_d := \mathcal{S}(\mathbb{R}^d) \subset L^2_d$  and is flanked by the topology induced by the countable family of seminorms  $\|\cdot\|_{n,k}$  which makes it a Fréchet space (so complete and locally convex  $\mathbb{T}_2$  topological metrizable vector space). A *tempered or slowly increasing distribution* on  $\mathbb{R}^d$  is a function from  $\mathcal{S}_d$  into  $\mathbb{R}$  which is linear and (sequentially) continuous w.r.t. that topology on  $\mathcal{S}_d$ . The space of the tempered distributions on  $\mathbb{R}^d$  is then the dual space  $\mathcal{S}'_d := (\mathcal{S}_d)' \subset \mathcal{D}'_d$  of  $\mathcal{S}_d$  and is coupled with the weak-star topology. As far as  $d = 1$ , we omit the subscript  $d$  from each of the spaces above.

**Theorem** (Bochner-Minlos-Sazonov). *There exists a complete probability measure  $\mathbf{P}$  defined on the Borel  $\sigma$ -algebra  $\mathcal{F}$  on  $\Omega := \mathcal{S}'$  such that, for every  $\varphi \in \mathcal{S}$ ,*

$$\int_{\Omega} e^{i\langle \omega; \varphi \rangle} \mathbf{P}(d\omega) = \exp\left(-\frac{1}{2} \|\varphi\|_{L^2}^2\right).$$

We name  $\mathbf{P}$  the *white noise probability measure* and  $(\Omega, \mathcal{F}, \mathbf{P})$  the *white noise probability space*. The (*smoothed*) *white noise process* is the map  $w: L^2 \rightarrow L^2(\mathbf{P})$  identified by placing, for  $\varphi \in \mathcal{S}$  and  $\omega \in \Omega$ ,

$$w_\varphi(\omega) := \langle \omega; \varphi \rangle$$

and by using then the  $\|\cdot\|_{L^2}$ -density of  $\mathcal{S}$  in  $L^2$ . Note that  $w$  is a linear isometry between Hilbert spaces.

Starting from  $w$ , one could easily construct a 1D standard Brownian motion  $W = (W(t))_{t \in \mathbb{R}}$  on  $\Omega$  ( $\mathbf{P}$ -a.s. null for negative times) in such a way that, for any  $f \in L^2$ ,

$$w_f = \int_{\mathbb{R}} f(t) dW(t)$$

which is a Wiener-Itô integral on  $\mathbb{R}$ . Let's settle  $\mathbb{F} = (\mathcal{F}_t)_{t \in \mathbb{R}}$  as the filtration generated by  $W$ .

For  $n \in \mathbb{N}$ , let  $H_n \in \mathbb{R}[x; n]$  be the  $n$ -th Hermite polynomial (on  $\mathbb{R}$ ), which is the  $n$ -th coefficient in  $x \in \mathbb{R}$  of the series expansion in powers of  $t \in \mathbb{R}$  of the smooth function  $(t, x) \mapsto \exp(tx - t^2/2)$ , namely

$$H_n(x) = \frac{(-1)^n}{n!} e^{x^2/2} \frac{d^n}{dx^n} e^{-x^2/2}$$

and let  $(\varphi_n)_{n \in \mathbb{N}^*}$  be the orthonormal basis of  $L^2$  consisting of the Hermite functions: for any  $x \in \mathbb{R}$ ,

$$\varphi_n(x) = \pi^{-1/4} (n-1)!^{-1/2} e^{-x^2/2} H_{n-1}(\sqrt{2}x).$$

Indicate with  $\Lambda = (\Lambda, +)$  the space of the multi-indices, i.e. the sequences  $k = (k_n)_{n \in \mathbb{N}^*}$  in  $\mathbb{N}$  with  $k_n \neq 0$  only for a finite number  $\ell(k)$  of them, and by the way set:  $|k| = \sum_{n=1}^{\infty} k_n$ ,  $k! = \prod_{n=1}^{\infty} k_n!$ ,  $0 := (0)_{n \in \mathbb{N}^*}$  and, for  $j \in \mathbb{N}^*$ ,  $\delta_{(j)} := (\delta_{j,n})_{n \in \mathbb{N}^*}$ . Define, for every  $k = (k_n)_{n \in \mathbb{N}^*} \in \Lambda$ ,

$$\Phi_k := \sqrt{k!} \prod_{n=1}^{\infty} H_{k_n}(w_{\varphi_n})$$

and, for  $j \in \mathbb{N}^*$ ,  $\Phi_{(j)} := \Phi_{\delta_{(j)}} = w_{\varphi_j}$ . Then  $(\Phi_k)_{k \in \Lambda}$  is an orthonormal basis of  $L^2$  and, for any  $t \geq 0$ ,

$$W(t) = \sum_{j=1}^{\infty} \left( \int_0^t \varphi_j(r) dr \right) \Phi_{(j)}.$$

A *Hida test function* on  $\Omega$  is a random variable  $X = \sum_{k \in \Lambda} a_k \Phi_k \in L^2(\mathbf{P})$ , with  $(a_k)_{k \in \Lambda}$  in  $\mathbb{R}$ , such that if, for any  $k = (k_n)_{n \in \mathbb{N}^*} \in \Lambda$ ,  $(k_{(1)}, \dots, k_{(\ell(k))})$  denotes the ordered  $\ell(k)$ -tuple formed by the non-zero components of  $k$  then, for every  $p \in \mathbb{R}$ ,

$$\|X\|_{(p)}^2 := \sum_{k \in \Lambda} k! a_k^2 \prod_{n=1}^{\ell(k)} (2n)^{p k_{(n)}} < \infty.$$

The space of the Hida test functions on  $\Omega$  is the *Hida test function space*  $(\mathcal{S}) \subset L^2(\mathbf{P})$  and is equipped by the projective topology induced by the family of norms  $\|\cdot\|_{(p)}$ . For instance, for any  $t \in \mathbb{R}$ ,  $W(t) \in (\mathcal{S})$ .

A *Hida distribution* on  $\Omega$  is a formal series  $Y = \sum_{k \in \Lambda} b_k \Phi_k$ , with  $(b_k)_{k \in \Lambda}$  in  $\mathbb{R}$  and where  $\mathbf{E}[Y] := b_0$  is named the *generalized expectation* of  $Y$ , such that there exists  $q \in \mathbb{R}$  for which

$$\|Y\|_{(-q)}^2 := \sum_{k \in \Lambda} k! b_k^2 \prod_{n=1}^{\ell(k)} (2n)^{-q k_{(n)}} < \infty.$$

The space of the Hida distributions on  $\Omega$  is the *Hida distribution space* and results the dual space  $(\mathcal{S})' \supset L^2(\mathbf{P})$  of  $(\mathcal{S})$  with  $\langle Y; X \rangle = \sum_{k \in \Lambda} k! a_k b_k$ . Here the  $\omega$ -pointwise product does not make sense.

The *singular ( $t$ -pointwise) white noise* on  $\Omega$  is given by, for any  $t \in \mathbb{R}$ , the distributional derivative  $\dot{W}(t) \in (\mathcal{S})'$  taken in  $(\mathcal{S})'$  of  $W(t)$  and thus, equivalently,

$$\dot{W}(t) = \sum_{j=1}^{\infty} \varphi_j(t) \Phi_{(j)}.$$

Well, for every  $X = \sum_{k \in \Lambda} a_k \Phi_k$  and  $Y = \sum_{k \in \Lambda} b_k \Phi_k$  in  $(\mathcal{S})'$ , the *Wick product*  $X \diamond Y$  of  $X$  and  $Y$  is the Hida distribution on  $\Omega$  defined as

$$X \diamond Y \doteq \sum_{k \in \Lambda} \left( \sum_{\alpha + \beta = k} a_\alpha b_\beta \right) \Phi_k.$$

The Wick algebra obeys the rules of an ordinary algebra, even together with the operation of sum, while caution is required about combining it with the ordinary product: in general, for arbitrary  $X, Y, Z \in (\mathcal{S})'$ ,  $X \cdot (Y \diamond Z) \neq (X \cdot Y) \diamond Z$  (whenever those pointwise products do make sense). Note also that in general, for arbitrary  $X, Y \in L^2(\mathbf{P})$ ,  $X \diamond Y \notin L^2(\mathbf{P})$  but, if  $X, Y \in (\mathcal{S})$ , then  $X \diamond Y \in (\mathcal{S})$  as well.

A map  $Y: \mathbb{R}_t \rightarrow (\mathcal{S})'$  is said  $(\mathcal{S})'$ -*integrable* if, for every  $X \in (\mathcal{S})$ ,  $\langle Y(\cdot); X \rangle \in L^1 := L^1(\mathbb{R}; \mathbb{R})$  and, in such a case, there exists an unique Hida distribution on  $\Omega$ , the  $(\mathcal{S})'$ -*integral*  $\int_{\mathbb{R}} Y(t) dt$  of  $Y$ , with

$$\left\langle \int_{\mathbb{R}} Y(t) dt; X \right\rangle = \int_{\mathbb{R}} \langle Y(t); X \rangle dt.$$

We remark that, if  $Y(\cdot)$  is  $(\mathcal{S})'$ -integrable then, for any  $T \in \mathbb{R}_+^*$ ,  $Y(\cdot) \mathbb{1}_{[0, T]}(\cdot)$  remains  $(\mathcal{S})'$ -integrable and we write  $\int_0^T Y(t) dt := \int_{\mathbb{R}} (Y(t) \mathbb{1}_{[0, T]}(t)) dt$ .

**Theorem.** Fix  $T \in \mathbb{R}_+^*$  and take  $u = (u(t))_{t=0}^T \in \text{Dom } \delta$  as a process on  $\Omega$ . Then, for any  $t \in [0, T]$ ,  $u(t) \diamond \dot{W}(t)$  is  $(\mathcal{S})'$ -integrable and its  $(\mathcal{S})'$ -integral coincides with the Skorohod integral of  $u$  (w.r.t.  $W$ ):

$$\int_0^T u(t) \diamond \dot{W}(t) dt = \int_0^T u(t) \delta W(t) \in L^2(\mathbf{P}).$$

For any map  $Y: \mathbb{R}_t \rightarrow (\mathcal{S})'$  such for which  $Y(\cdot) \diamond \dot{W}(\cdot)$  is  $(\mathcal{S})'$ -integrable,  $\int_{\mathbb{R}} Y(t) \diamond \dot{W}(t) dt$  is the *generalized Skorohod integral* of  $Y$ .

An entire construction like all that could be made in a very more general way and furthermore, anyhow, it could be genuinely connected to the Wiener-Itô chaos expansion.

## Rough volatility

Let  $(\Omega, \mathcal{F}, \mathbf{P})$  be a complete probability space such that there exists a 2D standard Brownian motion  $(W, \overline{W}) = ((W(t), \overline{W}(t)))_{t \in \mathbb{R}}$  on it ( $\mathbf{P}$ -a.s. null for negative times) and fix  $T \in \mathbb{R}_+^*$ .

On the one hand put  $\rho \in [-1, 1] \setminus \{0\}$ ,  $\bar{\rho} := \sqrt{1 - \rho^2}$  and produce the 1D standard Brownian motion  $B = (B(t))_{t \in \mathbb{R}}$  on  $\Omega$  defining its trajectories, for any  $t \in \mathbb{R}$ , as  $B(t) := \rho W(t) + \bar{\rho} \overline{W}(t)$  in such a way that  $B$  and  $W$  have constant correlation  $\rho \neq 0$ . Assign to  $\Omega$  the filtration  $\mathbb{F} = (\mathcal{F}_t)_{t \in \mathbb{R}}$  generated by  $B$ .

On the other hand choose a Hurst index  $H \in ]0, 1/2[$ , take the Volterra kernel  $K$  on  $\mathbb{R}_+^2$  given by, for any  $s, t \geq 0$ ,  $K(s, t) := K(t - s)$  where, for any  $r \in \mathbb{R}$ ,

$$K(r) := \sqrt{2H} |r|^{H-1/2} \mathbb{1}_{\mathbb{R}_+}(r)$$

and consider the 1D fractional Riemann-Liouville Brownian motion  $\widehat{W} = (\widehat{W}(t))_{t \in \mathbb{R}}$  on  $\Omega$  of index  $H$  and Volterra dynamics based on  $W$  having trajectories, for any  $t \in \mathbb{R}$ ,

$$\widehat{W}(t) := \int_0^t K(t-s) dW(s).$$

We remark that  $\widehat{W}$  is a continuous Gaussian process with negatively correlated increments and locally  $H^-$ -Hölder continuous trajectories, i.e. of any order strictly smaller than  $H$  and thus rougher than Brownian paths, which is a local martingale independent of  $\overline{W}$  that admits quadratic variation (by virtue of the classical Burkholder-Davis-Gundy inequality).

Fixed  $f \in C^1(\mathbb{R}; \mathbb{R}_+)$ , a (*simple*) *rough volatility* process  $\sigma = (\sigma(t))_{t \geq 0}$  on  $\Omega$  is explicitly given by

$$\sigma(t) := f(\widehat{W}(t)), \quad t \geq 0$$

meaning it indeed as the volatility process corresponding to the stochastic volatility Itô model

$$\begin{cases} dS(t)/S(t) = \sigma(t) dB(t), & t \geq 0, \\ S(0) \neq 0 & [\mathbf{P}]. \end{cases}$$

We're in the presence of a singular SDE due to the roughness of  $\sigma$  in the sense that, as  $\sigma$  is not even a semi-martingale, in particular it admits no Stratonovich form, closely related to which the absence of Markovianity of the model – although  $S = (S(t))_{t \geq 0}$  remains a local martingale – and the lack of a Wong-Zakai type approximation theory for that (and consequently the loss of hope in a successful use of the well-known tools and methods for SDEs).

We submit hereunder a regularity structure for  $\sigma$  which would be the basis for solving the above issues basically providing an approximation theory for stochastic integrals of type

$$\int f(\widehat{W}) dW.$$

Our task is to build an analysis à la Hairer based on renormalized enhanced noise, incorporating and keeping track of the relevant things we've in mind, so dealing with a mix of even hard algebraic-analytical conditions and, as usual in this area, with the problem of discovering the “right” approximation of the noise and therefore the renormalized approximating models.

Well, employed  $\kappa \in ]0, H[$  and  $M = M(H, \kappa) := \max \{m \in \mathbb{N} \mid m(H - \kappa) - 1/2 - \kappa \leq 0\} \in \mathbb{N}^*$ , define, always depending on  $M$ , the index set

$$A := \{-1/2 - \kappa, (H - \kappa) - 1/2 - \kappa, \dots, M(H - \kappa) - 1/2 - \kappa, 0, H - \kappa, \dots, M(H - \kappa)\}$$

and the symbols set

$$S := \{\Xi, \Xi \mathcal{I}(\Xi), \dots, \Xi \mathcal{I}(\Xi)^M, \mathbf{1}, \mathcal{I}(\Xi), \dots, \mathcal{I}(\Xi)^M\}$$

attaching to every symbol  $\tau$  in  $S$  a homogeneity  $|\tau|$  in  $A$  as follows:  $|\mathbf{1}| := 0$ ,  $|\Xi| := -1/2 - \kappa$  and, for  $m = 1, \dots, M$ ,  $|\Xi \mathcal{I}(\Xi)^m| := m(H - \kappa) - 1/2 - \kappa$  and  $|\mathcal{I}(\Xi)^m| := m(H - \kappa)$ . By reading powers of symbols as products with themselves, we see that homogeneities are multiplicative.

Of course  $\Xi$  should be interpreted as an abstract representation of the white noise  $\xi$  belonging to  $W$ , that is  $\xi = \dot{W}$  in the distributional sense; while  $\mathcal{I}(\cdot)$  has the intuitive meaning of integration against the Volterra kernel, improperly speaking, and in particular  $\mathcal{I}(\Xi)$  would perform  $\widehat{W}$ .

So let  $\mathcal{T} := \bigoplus_{\tau \in S} \langle \tau \rangle$  be the model space and  $G := \{ \Gamma_h \mid h \in (\mathbb{R}, +) \}$  be the structure group where

$$\Gamma_h \mathbf{1} := \mathbf{1}, \quad \Gamma_h \Xi := \Xi, \quad \Gamma_h \mathcal{I}(\Xi) := \mathcal{I}(\Xi) + h \mathbf{1} \quad \text{and} \quad \Gamma_h \tau \cdot \tau' := \Gamma_h \tau \cdot \Gamma_h \tau' \quad \text{for } \tau, \tau' \in S \text{ with } \tau \cdot \tau' \in S$$

then extending to  $\mathcal{T}$  by linearity and getting that the triple  $\mathcal{S} = (A, \mathcal{T}, G)$  is a regularity structure (also thanks to the basic binomial theorem which will serve throughout the continuation).

With the aim of building an appropriate limiting Itô model  $M = (\Pi, \Gamma)$  for  $\mathcal{S}$  (on  $\mathbb{R}$ ) based on  $\xi$ , for  $m = 1, \dots, M$  we give a meaning to the terms  $\Xi \mathcal{I}(\Xi)^m$  by defining an iterated Itô integral  $\mathbb{W}^m$  of second order so that it constitutes somehow a kind of its primitive (the context, for the moment, is too irregular to simply propose a product): for  $s, t \in \mathbb{R}$  with  $s \leq t$ ,

$$\mathbb{W}^m(s, t) := \int_s^t \left( \widehat{W}(r) - \widehat{W}(s) \right)^m dW(r).$$

By observing that this process satisfies  $\mathbf{P}$ -a.s. the pseudo Chen's relation, for  $s, u, t \in \mathbb{R}$  with  $s \leq u \leq t$ ,

$$\mathbb{W}^m(s, t) = \mathbb{W}^m(s, u) + \sum_{l=0}^m \binom{m}{l} \left( \widehat{W}(u) - \widehat{W}(s) \right)^l \mathbb{W}^{m-l}(u, t)$$

we extend  $\mathbb{W}^m$  to  $\mathbb{R}^2$  just by imposing that useful relation for every  $s, u, t \in \mathbb{R}$ : for  $s, t \in \mathbb{R}$  with  $t < s$ ,

$$\mathbb{W}^m(s, t) := - \sum_{l=0}^m \binom{m}{l} \left( \widehat{W}(t) - \widehat{W}(s) \right)^l \mathbb{W}^{m-l}(t, s).$$

**Lemma.** *For  $m = 1, \dots, M$  there exists a version of  $\mathbb{W}^m$ , for which we keep its symbol, and there exists  $p \in [1, \infty[$  such that, for any  $K \in \mathcal{K}$ , there exists a random positive  $t$ -constant  $C_K \in L^p(\mathbf{P})$  such that, for any  $s, t \in K$  (and  $\mathbf{P}$ -a.s.),*

$$|\mathbb{W}^m(s, t)| \leq C_K |s - t|^{m(H-\kappa)+1/2-\kappa}.$$

For  $s, t \in \mathbb{R}$  and  $m = 1, \dots, M$  (and  $\mathbf{P}$ -a.s.), define

$$\left\{ \begin{array}{l} \Gamma_{t,s} \mathbf{1} := \mathbf{1} \\ \Gamma_{t,s} \Xi := \Xi \\ \Gamma_{t,s} \mathcal{I}(\Xi) := \mathcal{I}(\Xi) + \left( \widehat{W}(t) - \widehat{W}(s) \right) \mathbf{1} \\ \Gamma_{t,s} \tau \cdot \tau' := \Gamma_{t,s} \tau \cdot \Gamma_{t,s} \tau' \quad \text{when } \tau \cdot \tau' \in S \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \Pi_s \mathbf{1} := \mathbf{1} \\ \Pi_s \Xi := \dot{W} \\ \Pi_s \mathcal{I}(\Xi)^m := \left( \widehat{W}(\cdot) - \widehat{W}(s) \right)^m \\ \Pi_s \Xi \mathcal{I}(\Xi)^m := \frac{d}{dt} \mathbb{W}^m(s, \cdot) \end{array} \right.$$

then extending to  $\mathcal{T}$  by linearity.

**Proposition.** *The pair  $M = (\Pi, \Gamma)$ , where  $\Gamma = (\Gamma_{t,s})_{t,s \in \mathbb{R}}$  and  $\Pi = (\Pi_s)_{s \in \mathbb{R}}$ , is ( $\mathbf{P}$ -a.s.) a model for  $\mathcal{S}$ .*

The next goal is to find, for  $\varepsilon \downarrow 0$ , a reasonable approximating model  $M^\varepsilon = (\Pi^\varepsilon, \Gamma^\varepsilon)$  for  $\mathcal{S}$  of  $M$ , to be renormalized, relying on a smart approximation  $\dot{W}^\varepsilon$  of  $\xi$  (as distributions). About this, let's consider a function  $\delta^\varepsilon: \mathbb{R}_{(x,y)}^2 \rightarrow \mathbb{R}$ , which should be understood as an approximation of the Dirac delta and which could be easily built from a mollifier as well as a wavelet basis, with the following properties.

- ★ The function  $\delta^\varepsilon$  is measurable, symmetric and bounded with  $\|\delta^\varepsilon\|_\infty = \mathcal{O}(\varepsilon^{-1})$ .
- ★ There exists  $\beta \in ]1/2 + \kappa, \infty[$  such that, for any  $x \in \mathbb{R}$ , the function  $\delta^\varepsilon(x, \cdot): \mathbb{R}_y \rightarrow \mathbb{R}$  belongs to the Besov space  $\mathcal{B}_{1,\infty}^\beta(\mathbb{R})$  and  $x \mapsto \delta^\varepsilon(x, \cdot)$  is measurable and bounded as a map from  $\mathbb{R}_x$  into  $\mathcal{B}_{1,\infty}^\beta(\mathbb{R})$ .
- ★ There exists  $c \in ]0, \infty[$  such that, for any  $x \in \mathbb{R}$ ,  $\text{supp } \delta^\varepsilon(x, \cdot) \subset B_{c\varepsilon}(x)$  with  $\int_{\mathbb{R}} \delta^\varepsilon(x, y) dy = 1$ .

Indeed  $\dot{W}$  turns out to be locally contained in  $\mathcal{B}_{\infty, \infty}^{-1/2-\kappa}(\mathbb{R}) \subset (\mathcal{B}_{1, \infty}^{\beta}(\mathbb{R}))'$  and therefore we define the approximation  $\dot{W}^{\varepsilon} := (\dot{W}^{\varepsilon}(t))_{t \in \mathbb{R}}$  of  $\xi$ , a Gaussian pathwise measurable locally bounded process, as

$$\dot{W}^{\varepsilon}(t) := \left\langle \dot{W}; \delta^{\varepsilon}(t, \cdot) \right\rangle \mathbb{1}_{\mathbb{R}_+}(t).$$

For 1D stochastic process  $u = (u(t))_{t=0}^T$  on  $\Omega$  and  $t \in \mathbb{R}_+$ , we write

$$\int_0^t u(r) dW^{\varepsilon}(r) := \int_0^t u(r) \dot{W}^{\varepsilon}(r) dr$$

while, if  $u$  takes values in some non-homogeneous *Wiener chaos* induced by  $\dot{W}$ , we write

$$\int_0^t u(r) \diamond dW^{\varepsilon}(r) := \int_0^t u(r) \diamond \dot{W}^{\varepsilon}(r) dr.$$

In particular, we consider the approximation  $\widehat{W}^{\varepsilon} := (\widehat{W}^{\varepsilon}(t))_{t \in \mathbb{R}}$  of  $\widehat{W}^{\varepsilon}$  as

$$\widehat{W}^{\varepsilon}(t) := K * \dot{W}^{\varepsilon} = \int_0^t K(t-r) dW^{\varepsilon}(r).$$

**Lemma.** For  $\varepsilon \downarrow 0$ , there exist  $p \in [1, \infty[$  and random positive  $t$ -constants  $C_{\varepsilon, T}, C_T \in L^p(\mathbf{P})$ , where are uniformly bounded, such that, for any  $s, t \in [0, T]$ ,  $\kappa' \in ]0, H[$  and  $\delta \in ]0, 1[$  (and  $\mathbf{P}$ -a.s.),

$$\left| \widehat{W}^{\varepsilon}(t) - \widehat{W}^{\varepsilon}(s) \right| \leq C_{\varepsilon, T} |t-s|^{H-\kappa'} \quad \text{and} \quad \left| \widehat{W}^{\varepsilon}(t) - \widehat{W}^{\varepsilon}(s) - \left( \widehat{W}(t) - \widehat{W}(s) \right) \right| \leq C_T |t-s|^{H-\kappa'} \varepsilon^{\delta \kappa'}.$$

For  $\varepsilon \downarrow 0$ ,  $s, t \in \mathbb{R}$  and  $m = 1, \dots, M$  (and  $\mathbf{P}$ -a.s.), define the approximating model for  $\mathcal{I}$  of  $M$  by

$$\left\{ \begin{array}{l} \Gamma_{t,s}^{\varepsilon} \mathbf{1} := \mathbf{1} \\ \Gamma_{t,s}^{\varepsilon} \Xi := \Xi \\ \Gamma_{t,s}^{\varepsilon} \mathcal{I}(\Xi) := \mathcal{I}(\Xi) + (\widehat{W}^{\varepsilon}(t) - \widehat{W}^{\varepsilon}(s)) \mathbf{1} \\ \Gamma_{t,s}^{\varepsilon} \boldsymbol{\tau} \cdot \boldsymbol{\tau}' := \Gamma_{t,s}^{\varepsilon} \boldsymbol{\tau} \cdot \Gamma_{t,s}^{\varepsilon} \boldsymbol{\tau}' \text{ when } \boldsymbol{\tau} \cdot \boldsymbol{\tau}' \in S \end{array} \right. \quad \text{and} \quad \left\{ \begin{array}{l} \Pi_s^{\varepsilon} \mathbf{1} := \mathbf{1} \\ \Pi_s^{\varepsilon} \Xi := \dot{W}^{\varepsilon} \\ \Pi_s^{\varepsilon} \mathcal{I}(\Xi)^m := (\widehat{W}^{\varepsilon}(\cdot) - \widehat{W}^{\varepsilon}(s))^m \\ \Pi_s^{\varepsilon} \Xi \mathcal{I}(\Xi)^m := \dot{W}^{\varepsilon}(\cdot) (\widehat{W}^{\varepsilon}(\cdot) - \widehat{W}^{\varepsilon}(s))^m \end{array} \right.$$

(extending then to  $\mathcal{T}$  by linearity).

**Proposition.** The pair  $M^{\varepsilon} = (\Pi^{\varepsilon}, \Gamma^{\varepsilon})$ , where  $\Gamma^{\varepsilon} = (\Gamma_{t,s}^{\varepsilon})_{t,s \in \mathbb{R}}$  and  $\Pi^{\varepsilon} = (\Pi_s^{\varepsilon})_{s \in \mathbb{R}}$ , is a model for  $\mathcal{I}$ .

In order to understand the diverging quantities  $\mathcal{C}^{\varepsilon}$  to be subtracted from  $\Pi^{\varepsilon}$  to renormalize it in the sense of Hairer, since there cannot be any hope of a convergence like  $\int f(\widehat{W}^{\varepsilon}) dW^{\varepsilon}$  to  $\int f(\widehat{W}) dW$ , what's below is enlightening. Before, given 1D Gaussians  $U_1, V, U_2$  on  $\Omega$  and  $l, h \in \mathbb{N}^*$  with  $l \leq h$ ,

$$U_1^l \cdot (V \diamond U_2^{h-l}) = V \diamond (U_1^l U_2^{h-l}) + l \mathbf{E}[V U_1] U_1^{l-1} U_2^{h-l}.$$

**Lemma.** For any  $\varphi \in \mathcal{D}$ ,  $s \in \mathbb{R}$ ,  $m = 1, \dots, M$  and  $\varepsilon \downarrow 0$ , the two following identities hold.

$$1. \Pi_s \Xi \mathcal{I}(\Xi)^m(\varphi) = \int_0^{\infty} \varphi(t) (\widehat{W}(t) - \widehat{W}(s))^m \delta W(t) - m \int_0^s \varphi(s) K(s-r) (\widehat{W}(r) - \widehat{W}(s))^{m-1} dr.$$

$$2. \text{Defined } \mathcal{K}^{\varepsilon}(s, \cdot) := \mathbf{E}[\widehat{W}^{\varepsilon}(s) \dot{W}^{\varepsilon}(\cdot)] = \mathbb{1}_{\mathbb{R}_+^2}(s, \cdot) \int_0^{\infty} \int_0^{\infty} \delta^{\varepsilon}(\cdot, x) \delta^{\varepsilon}(x, y) K(s-y) dx dy, \text{ then}$$

$$\begin{aligned} \Pi_s^{\varepsilon} \Xi \mathcal{I}(\Xi)^m(\varphi) &= \int_0^{\infty} \varphi(t) (\widehat{W}^{\varepsilon}(t) - \widehat{W}^{\varepsilon}(s))^m \diamond dW^{\varepsilon}(t) \\ &\quad - m \int_0^{\infty} \varphi(t) [\mathcal{K}^{\varepsilon}(s, t) - \mathcal{K}^{\varepsilon}(t, t)] (\widehat{W}^{\varepsilon}(t) - \widehat{W}^{\varepsilon}(s))^{m-1} dt \end{aligned}$$

Furthermore, assuming  $s \leq T$ , there exists  $C_T \in \mathbb{R}_+^*$  such that, for any  $t \in [0, T]$ ,

$$|\mathcal{K}^{\varepsilon}(s, t)| \leq C_T \varepsilon^{H-1/2}.$$



As a corollary, if we interpret  $\mathcal{K}^\varepsilon$  as an approximation of the kernel  $K$ , then we see that

$$\mathcal{C}^\varepsilon(t) := \mathcal{K}^\varepsilon(t, t), \quad t \geq 0$$

would correspond to something diverging like “ $0^{H-1/2} = \infty$ ” in the limit  $\varepsilon \downarrow 0$ .

**Theorem.** For  $\varepsilon \downarrow 0$ ,  $s \in \mathbb{R}$  and  $m = 1, \dots, M$  (and  $\mathbf{P}$ -a.s.), define

$$\widehat{\Pi}_s^\varepsilon \Xi \mathcal{I}(\Xi)^m := \Pi_s^\varepsilon \Xi \mathcal{I}(\Xi)^m - m \mathcal{C}^\varepsilon(\cdot) \Pi_s^\varepsilon \mathcal{I}(\Xi)^{m-1}$$

leaving  $\widehat{\Pi}_s^\varepsilon \boldsymbol{\tau} := \Pi_s^\varepsilon \boldsymbol{\tau}$  on the remaining symbols  $\boldsymbol{\tau} \in S$ . Then the pair  $\widehat{M}^\varepsilon = (\widehat{\Pi}^\varepsilon, \Gamma^\varepsilon)$ , where  $\widehat{\Pi}^\varepsilon = (\widehat{\Pi}_s^\varepsilon)_{s=0}^T$ , is a model for  $\mathcal{S}$  and there exists  $C_T \in \mathbb{R}_+^*$  such that, for any  $p \in [1, \infty[$  and  $\delta \in ]0, 1[$ ,

$$\left\| \left\| \widehat{M}^\varepsilon; \mathbf{M} \right\| \right\|_T \Big\|_{L^p(\mathbf{P})} \leq C_T \varepsilon^{\delta \kappa}$$

where of course, for any model  $\widetilde{M} = (\widetilde{\Pi}, \widetilde{\Gamma})$  for  $\mathcal{S}$  on  $\mathbb{R}$ ,

$$\begin{aligned} \left\| \left\| \widetilde{M}; \mathbf{M} \right\| \right\|_T \doteq & \sup \left\{ \left| (\Pi_s - \widetilde{\Pi}_s) \boldsymbol{\tau}(\varphi_s^\lambda) \right| \lambda^{-|\boldsymbol{\tau}|} \mid \varphi \in \mathcal{D}_1, \lambda \in ]0, 1], s \in [0, T], \boldsymbol{\tau} \in S \right\} \\ & + \sup \left\{ \left\| \Gamma_{t,s} \boldsymbol{\tau} - \widetilde{\Gamma}_{t,s} \boldsymbol{\tau} \right\|_{\alpha'} \mid |t - s|^{\alpha' - |\boldsymbol{\tau}|} \mid t, s \in [0, T], \boldsymbol{\tau} \in S, \alpha' \in A \text{ with } \alpha' < |\boldsymbol{\tau}| \right\}. \end{aligned}$$

---

### Starting reference

[1] C. Bayer, P. K. Friz, P. Gassiat, J. Martin, B. Stemper. *A regularity structure for rough volatility*. *Mathematical Finance* (2019), pp. 1–51.